

RC
114.5
R943
1961

Bethesda

UNCLASSIFIED

AD 261 142

*Reproduced
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

261142

CATALOGED BY ASTIA

AS AD No


TECHNICAL STUDY 33

RIFT VALLEY FEVER A REVIEW OF THE LITERATURE

1960-62

JULY 1961

XEROX



U.S. ARMY CHEMICAL CORPS
BIOLOGICAL LABORATORIES
FORT DETRICK

Copy 57 of 88 copies

U.S. ARMY CHEMICAL CORPS RESEARCH AND DEVELOPMENT COMMAND
U.S. ARMY BIOLOGICAL LABORATORIES
Fort Detrick, Maryland

Technical Study 33

RIFT VALLEY FEVER

A Review of the Literature

Johnnie L. Runnels

"

Leslie C. Murphy

Virus and Rickettsia Division
DIRECTOR OF BIOLOGICAL RESEARCH

Project 4B11-02-065

July 1961

XC114.5
R943
1961

This document or any portion thereof may not be reproduced without specific authorization from the Commanding Officer, Biological Laboratories, Fort Detrick, Frederick, Maryland; however, ASTIA is authorized to reproduce the document for U.S. Government purposes.

The information in this report has not been cleared for release to the general public.

ASTIA AVAILABILITY NOTICE

Qualified requestors may obtain copies of this document from ASTIA.

Foreign announcement and dissemination of this document by ASTIA is limited.

FOREWORD

The literature pertaining to Rift Valley fever has been reviewed in conjunction with investigation of the disease and its etiologic virus. The work was performed under Project 4B11-02-065.

Intended as an economic measure, it has been addressed to the consolidation and summarization of published material of potential value in the study, research, and evaluation of Rift Valley fever.

The number of reports pertaining to this disease and its agent was limited by the number of laboratories engaged in investigations of the disease since its discovery in 1931. Approximately 180 reports were reviewed. The number of reports published world-wide was estimated at about 200. Because of duplication only about 100 of the available reports were cited in this text.

THE UNIVERSITY OF CHICAGO
LIBRARY
1000 1000

1000 1000
1000 1000
1000 1000

1000 1000
1000 1000
1000 1000
1000 1000

4

DIGEST

Rift Valley fever literature was reviewed with the intention of accumulating, under a single cover, published material of use in the study, research, and evaluation of the disease. Emphasis was placed on presentation of facts as they were reported in scientific publications.

Attention has been given to (a) geographic distribution, (b) modes of transmission, (c) susceptibility of hosts, (d) pathology, (e) immunity, (f) characteristics of the virus, and (g) investigation procedures.

Rift Valley fever has been shown to be a highly infectious disease of sheep, cattle, and other animals. Man has been infected frequently during epizootics of domestic animals and during laboratory or field contact with infectious material. The disease has not been found to occur naturally outside the African continent, but accidental human infections have been reported in the United States, Europe, and Japan.

CONTENTS

Foreword	3
Digest	4
I. INTRODUCTION	7
A. Definition	7
B. History	7
II. RIFT VALLEY FEVER	8
A. Geographic Distribution	8
B. Susceptibility	10
C. Transmission	10
1. Natural	10
2. Experimental	11
3. Accidental	12
D. Human Infections	13
1. Sources	13
2. Experimental Infection	13
3. Confirmed Infections	14
4. Morbidity and Mortality	14
5. Clinical Picture	14
6. Immunity	19
E. Animal Infections	20
1. Natural Infections	20
2. Experimental Infections	21
3. Pathology	21
4. Immunization	24
F. Diagnosis	25
G. Treatment	25
III. RIFT VALLEY FEVER VIRUS	26
A. Recognized Strains	26
1. Pantropic	26
2. Neurotropic	26
3. Lunyo Virus	27
B. Morphology	27
C. Propagation	28
1. Mice	28
2. Lambs	28
3. Embryonated Eggs	28
4. Tissue Culture	29
5. Adsorption and Multiplication	29
D. Stability	31
1. Acid Tolerance	31
2. Heat	32
3. Chemicals	32
4. Storage	32

E.	Identification Procedure	33
1.	Isolation	33
2.	Identification	34
F.	Classification	35
IV.	SUMMARY	36
	Literature Cited	37
	References	45

FIGURES

1.	Geographic Distribution of Rift Valley Fever in Africa	9
2.	Clinical Signs of Rift Valley Fever Infection in Humans	16

TABLES

I.	Rift Valley Fever Isolations from Wild-Caught Mosquitoes	11
II.	Vectors With Which Experimental Transmission of Rift Valley Fever Virus has been Achieved	11
III.	Mosquitoes that were Infected with Rift Valley Fever Under Laboratory Conditions but did not Transmit Virus	12
IV.	Summary of Rift Valley Fever Infections in Humans	15
V.	Ocular Sequelae in Human Cases of Rift Valley Fever	18
VI.	Circulation of PRVFEV in Blood of Various Hosts	34

I. INTRODUCTION

A. DEFINITION

Rift Valley fever has been defined as an acute, febrile, insect-borne virus disease of sheep, cattle, and other animals. Also susceptible, man has been infected during epizootics of domestic animals and laboratory accidents.

B. HISTORY

In 1930, Daubney et al.^{1/*} isolated a specific virus of the disease during an epizootic in sheep and cattle of the Rift Valley in Africa. The Rift Valley was described as a huge geological depression that starts in Persia and continues through northeastern and central Africa until it ends in eastern Transvaal.^{2/} However, the disease was not confined to the Rift Valley, and it probably existed in Africa much earlier than 1931. According to Weiss,^{3/} Montgomery (1912) and Stordy (1913) reported the existence of a disease in man and domestic animals with a symptomatic resemblance to Rift Valley fever.

Daubney first designated the disease "enzootic hepatitis" because of extensive liver damage in infected animals. Because pathology extended to other organs as well, he later decided upon the name Rift Valley fever, the designation consistently employed in the literature.^{1/}

* See Literature Cited.

II. RIFT VALLEY FEVER

A. GEOGRAPHIC DISTRIBUTION

Evidence of Rift Valley fever has been found in a wide belt extending from the southern tip of the Union of South Africa northward into Central Africa.^{4/} Occurrences of the disease in this belt and elsewhere are shown in Figure 1.

The first proved epizootic, described by Daubney in 1931, occurred in Kenya.^{1/} The disease was contracted by shepherds attending infected flocks and by four Europeans engaged in examination of infected animals and in laboratory work with infected materials.

Transmission of the virus by mosquitoes and presence of the disease in Uganda was reported by Smithburn *et al.* in 1948, when virus was isolated from mosquitoes caught in the uninhabited Semliki forest.^{5/} Even before this, Findlay *et al.*^{6/} had shown that virus-neutralizing antibodies existed in the sera of natives in Uganda, French Sudan, Anglo-Egyptian Sudan, and French Equatorial Africa. Another outbreak of the disease occurred in Kenya in 1933.^{7/} In the summers of 1950 and 1951 the first epizootic occurred in the Union of South Africa.^{8,9/} The South Africa epizootic involved the western and southwestern areas of Orange Free State, northern and western areas of Cape Province, and the western and southern areas of Transvaal.^{10/} Epizootics reappeared in the Union of South Africa in 1953 (Fauresmith district), 1955 (Mariesburg district), and in the summer of 1956 (Western Orange Free State).^{3,11-13/} In 1957 Kokernot *et al.*^{14/} found that wild-caught mosquitoes in Zululand carried the virus and Jhone reported the presence of antibodies in the sera of cattle in Southern Rhodesia.^{15/}

Kaschula^{17/} found antibodies in the sera of cattle in the Knysna district, but no evidence of clinical disease other than an occasional abortion in cattle. He suggested that these areas formed truly enzootic areas similar to the Semliki forest. Enzootic areas were described as low, warm areas with high annual rainfall. Similar terrain and climate plus the added factor of sheep farming characterized the epizootic areas. It was reasoned that the disease reached epizootic proportions only in those enzootic areas that were also suitable for sheep, a highly susceptible host.^{3,18/} The disease halted abruptly with the first frost and with movement of herds to high altitudes.^{1,7/} The geographic distribution of arthropod-borne viruses in South Africa was investigated in some detail by Kokernot *et al.*^{14,18-20/} and Paterson *et al.*^{21/}




Figure 1. Geographic Distribution of Rift Valley Fever in Africa.

B. SUSCEPTIBILITY

A wide range of species was classified according to their susceptibility to laboratory infection with Rift Valley fever by Findlay *et al.*^{22-24/} He classified lambs, mice, hamsters, and wild rodents as most susceptible, inasmuch as the disease was usually fatal to these animals. It was also fatal to about fifty per cent of the second group, composed of sheep and rats. Men, monkeys, cows, goats, and grey squirrels experienced severe nonfatal infections and were placed in a third group. Cats were next with a mild reaction, followed by rabbits, which showed no evidence of infection although the virus persisted in the blood for several days. In a study of susceptibility among various species of monkeys, Findlay reported differences in susceptibility that were separate from differences in antibody response.^{25/} Indian, African, and South American monkeys were equally susceptible, although certain species of African monkeys were less susceptible. The more susceptible group exhibited a febrile reaction and viremia in contrast to certain African monkeys, which showed nothing more than the circulation of virus for a few days. In 1952 Findlay encountered differences in the susceptibility of mice and rats in relation to age and diet.^{26/} Suckling mice were more susceptible than older mice and rats. Rats maintained on an inadequate diet were more susceptible than well-fed rats. Findlay pointed out that the diet upon which the rats were maintained for the susceptibility studies in 1932 was not considered adequate in the 1952 studies. High death rates indicated that ferrets should be placed in the group of highly susceptible animals.^{27/} Horses, pigs, mongooses, hedgehogs, tortoises, frogs, and domesticated and wild fowl were listed as nonsusceptible by Findlay.^{22/}

C. TRANSMISSION

1. Natural

Seasonal aspects of Rift Valley fever outbreaks coincident with prevalence in low terrain following periods of heavy rainfall led Daubney *et al.* to conclude that the virus was transmitted naturally by biting insects.^{1/} Furthermore, they were able to stop the disease by moving flocks to higher altitudes or sheltering animals from mosquitoes. The mosquito was established as a vector by Smithburn *et al.* in 1948 when the virus was isolated from mosquitoes caught in the Semliki forest of Uganda.^{5/} Virus was isolated from three species of the genus *Aedes* and six species of the genus *Eretmapodites*. Material from the *Aedes* species had lower titer than that from the *Eretmapodites* species. It was thus concluded that an *Eretmapodites* species was the vector and the *Aedes* species were incidentally infected and not involved in the parasitic cycle. In 1949 the virus was transmitted from lamb to lamb and from mouse to mouse in the laboratory with *Eretmapodites chrysogaster*.^{28/} Epidemiological studies in 1948, 1951, and 1953 showed no evidence of person-to-person spread or infection of man by an insect vector.^{5,12,29/} Virus isolations from wild-caught mosquitoes are summarized in Table I.




TABLE I. RIFT VALLEY FEVER ISOLATIONS FROM WILD-CAUGHT MOSQUITOES

SPECIES	LOCATION	DATE	LITERATURE CITED
<u>Aedes deboeri de-meilloni</u>	Africa	1948	5
<u>Aedes tarsalis</u>	Africa	1948	5
<u>Aedes caballus</u>	South Africa	1955	13
<u>Aedes circumluteolus</u>	South Africa	1956 & 1957	20,30
<u>Aedes africanus</u>	South Africa	1956 & 1958	30
<u>Eretmapodites chrysogaster</u>	Africa	1948	5
<u>Culex theileri</u>	South Africa	1955	13

2. Experimental

Rift Valley fever virus has been transmitted in the laboratory by scarification, injection, nasal instillation, eye instillation, injection by biting insects, and sometimes by eating infected material. Vectors that have been induced to transmit the virus in the laboratory and those that have become infected without transmitting the virus are shown in Tables II and III, respectively.

TABLE II. VECTORS WITH WHICH EXPERIMENTAL TRANSMISSION OF RIFT VALLEY FEVER VIRUS HAS BEEN ACHIEVED

SPECIES	DATE	LITERATURE CITED	REMARKS
<u>Rhipicephalus appendiculatus</u> (tick)	1933	7	Virus not retained thru molt to adult
<u>Eretmapodites chrysogaster</u>	1949	28	
<u>Aedes aegypti</u>	1955	31,32	Adapted viscerotropic Lunyo strain only
<u>Aedes caballus</u>	1955	13	
<u>Aedes tarsalis</u>	1954	2	
<u>Mansonia</u> species	1954	2	

TABLE III. MOSQUITOES THAT WERE INFECTED WITH
RIFT VALLEY FEVER UNDER LABORATORY CONDITIONS
BUT DID NOT TRANSMIT VIRUS

SPECIES	DATE	LITERATURE CITED
<u>Culex theileri</u>	1955	33
<u>Mansonia fuscopennata</u>	1933	7
<u>Mansonia versicolor</u>	1933	7
<u>Mansonia microannulata</u>	1933	7

The immediate portal through which the virus infects during ingestion of food has not been clearly shown. Daubney *et al.* failed to transmit the virus by drenching a lamb with infected blood.^{1/} Yet mice have been infected by allowing them to feed on other mice that were moribund or dead of Rift Valley fever.^{32/} It was pointed out that actual transmission could have occurred in several ways. Experimental transmission of virus by nasal and eye instillation^{34/} and scarification suggested that infection could have resulted from cage-dust aerosol, aerosol created by clawing at infected mice, or scratches incurred during eating, but probably not from simple ingestion of infected material.^{22/} Experimental transmission of the virus to a human volunteer is described in the section on human infections.

3. Accidental

Exact portals of entry responsible for transmission of the virus in accidental infection of veterinarians and laboratory workers remain somewhat obscure. The first reported infections followed participation in post-mortem examination of infected animals and laboratory exposure in which little or no protective paraphernalia was used.^{1/} Later, varying degrees of protection including gloves, surgical masks, and protective over-garments were used without success. Accidental infection has also resulted from handling contaminated glassware,^{32/} entering rooms housing infected animals, pouring virus suspensions, and grinding infected tissue in mortar and pestle.^{35/} One accidental infection occurred under circumstances that suggested unusual resistance of the virus to drying.^{27/} An individual, whose presumed contact with the virus was scraping and painting the walls of a room that had housed infected laboratory animals three months earlier, became infected 15 days after the paint-scraping experience. Active work in that location with Rift Valley fever virus had been terminated four months before the onset of illness.

1. The first part of the document is a list of names and addresses, which appears to be a directory or a list of subscribers. The names are written in a cursive script, and the addresses are listed below them.

Findlay believed that the virus could gain entry through skin abrasions, the conjunctival sac, or the mucous membrane of the nose.^{22/}

Investigators have failed to isolate the virus from the urine of infected man or animals, but it has been isolated from muco-hemorrhagic feces^{36/} and material aborted by infected ewes.^{17/}

Search has been made for an intermediate host that circulated an adequate titer of virus long enough to participate in the parasitic cycle. Weinbren and Mason^{37/} observed that a wild field rat, Arvicanthis abyssinicus, had antibodies to Rift Valley fever in its serum and investigated the animal as a possible intermediate host. They concluded that the rodent could act as a natural host to the virus.

D. HUMAN INFECTIONS

1. Sources

Rift Valley fever infections acquired under natural conditions occurred in humans during each recorded epizootic of sheep and cattle.^{3/} Infections occurred among personnel engaged in post-mortem examination of diseased animals, among laboratory personnel in contact with infectious materials and among others whose occupations involved them with infected animals.^{2/}

2. Experimental Infection

Daubney et al.^{1/} provided the only report of a human experimental infection with Rift Valley fever virus. No apparent risk was involved, because earlier accidental infections were without known fatalities. An adult male of the Kissi tribe, a malarial patient, was given three cubic centimeters of diluted virus intramuscularly. The inoculum was prepared from 0.2 cc of lamb plasma filtrate which was obtained with a Chamberland L5 filter. Titration of the material was not reported, but the patient probably received 10^7 to 10^8 MIPLD₅₀ of virus.

On the third day after inoculation, the patient complained of headache and pain in the loins. On the fourth day, symptoms became severe and the febrile response appeared. The face was congested, eyes were slightly bloodshot, and the pulse was rapid. On the fifth day the temperature returned to normal but abdominal discomfort remained for weeks. Viremia was detected from the fourth through the ninth days after inoculation. No untoward sequelae were reported.

The above information was obtained from a review of the files of the Department of the Interior, Bureau of Land Management, and the National Archives and Records Administration.

1940

1994

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

116

1. 1993

1924

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 26

100

444

501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000

... ..

... ..

h. 72

3. Confirmed Infections

Rift Valley fever infections in humans have been heavily documented from case reports of persons contracting the disease during occupational contacts with the virus. Reports of confirmed cases have been summarized in Table IV. Note that the first such report was made by Daubney and co-workers in 1931. Infections were contracted in every laboratory in which work with Rift Valley fever was carried on. Infections of African, European, Japanese, and American workers in a variety of occupations were reported. Most infections followed known contact, but contact was not always well defined. High morbidity among those exposed to the virus was apparent from the earliest reports.

4. Morbidity and Mortality

Daubney *et al.*^{1/} and Findlay^{22/} reported 100 per cent morbidity among personnel following post-mortem experience with infected carcasses. Almost every native engaged in herding sheep during the 1930 epizootic contracted a disease with symptomatic resemblance to Rift Valley fever. Subsequent to the 1950 outbreak in South Africa, Schulz estimated 70 per cent morbidity among approximately 32,000 people assumed to be directly exposed.^{10/} Ten to fifteen per cent morbidity was estimated for the total population assumed to be at risk, approximately 500,000 people. The number of persons physically involved with infectious materials was much smaller. All state veterinarians, stock inspectors, and persons known to take an active part in post-mortem examination of diseased animals contracted the fever. Thus, the morbidity among susceptibles positively exposed was 100 per cent; lower occurrence in the general population probably reflected degree of exposure rather than morbidity.

In contrast to high morbidity, only one human fatality associated with Rift Valley fever has been reported.^{39/} This report is reviewed in Section II, D,5,d.

5. Clinical Picture

a. Symptoms

Remarkably similar symptomatically to dengue fever, Rift Valley fever has a sudden onset with elevated temperature, headache, muscular pain, weakness, sensation of fullness over the liver, rigors, vertigo, photophobia, nausea, and sometimes constipation and epistaxis.^{3/} Elevated temperature frequently occurs in two phases.^{1,22/}

b. Incubation Period

After three to six days' incubation period, a rise in temperature occurs, along with one or more of the symptoms previously described,

This is a report of the results of the investigation of the case of the missing person, Mr. John Doe, who was last seen on the 15th of March, 1964. The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

1. General Information

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

2. Details of the Case

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

The investigation was conducted by the police department of the city of New York, and the results are as follows:

TABLE IV. SUMMARY OF RIFT VALLEY FEVER INFECTIONS IN HUMANS


PLACE	NO. OF CASES AND OCCUPATION	CONFIRMATION TEST ^a	LITERATURE CITED
Kenya	2 veterinary surgeons 2 lab assistants	None	1
London	1 pathologist 1 veterinary surgeon 3 lab assistants	SN SN V, SN	22, 38
United States	1 pathologist 3 technicians 2 lab assistants 2 virologists	V, SN	27, 35, 38, 39
Uganda	2 pathologists 6 technicians 3 animal caretakers	V, SN	40
Japan	12 lab workers	CF, SN	41
South Africa	14 farmers 7 veterinary surgeons 2 teachers 2 research station employees 1 merchant 1 diamond worker 1 technician 2 natives 8 unknown	V, SN, CF	12, 15, 17, 42-47

- a. SN = serum neutralization
V = virus isolated
CF = complement fixation

and persists for two to three days (Figure 2). The temperature frequently returns to normal for one or two days and rises a second time. Temperatures reach 103° to 107°F, usually accompanied by symptoms.^{3/}

c. Incapacitation

The febrile response more or less paralleled the period during which virus was isolated from the blood, a period of four to six days.^{22/}



Adapted from Daubney *et al.*^{1/} and Findlay^{22/}

Figure 2. Clinical Signs of Rift Valley Fever Infection in Humans.

This period was also marked by a sharp rise in the leucocyte count, which changed to leucopenia as the febrile period terminated. Incapacitation varied from inapparent infections to complete debilitation. However, clinically recognizable cases usually required bed rest during the febrile period. Recovery was most often uninterrupted and complete, but malaise, weakness, and complaints of headache and defective vision were reported to persist for several weeks.^{3/}

d. Sequelae

(1) Ocular

Symptomatic visual disturbances were noted frequently in reports on human cases of Rift Valley fever. Photophobia, tenderness of the eyeballs, and pain behind the eyes were frequently reported.^{44/}

Daubney *et al.*^{1/} reported that a laboratory worker complained of defective vision for some weeks following the disease. The nature of the visual defect was not described, nor was there any reference to investigation of the complaint. Although complaints of visual disturbance have been made following Rift Valley fever infections in Central Africa, evidence of retinal damage following infection has been reported only in the Union of South Africa.^{32/}

Central serous retinopathy, characterized by macular swelling and occasional small hemorrhages, was described by Freed^{44/} and Schrire.^{45/} Freed reported the case of a schoolmaster, 38 years old, who complained of visual disturbance of six weeks' duration in the left eye. Six days after onset of illness the patient noticed blurring of vision. Six weeks later examination showed the cause to be a dense white elliptical mass covering the left macula. Previous exposure to Rift Valley fever virus was shown by the complement-fixation test.

Schrire^{45/} described five cases of macular exudates and one case of retinal detachment seen in his practice during the 1951 epizootic in the Union of South Africa. All of these cases were diagnosed by the complement-fixation test. Data pertinent to these cases are summarized in Table V. The onset of eye symptoms varied from the beginning of the febrile period to three weeks later. Visual defects remained in most of the cases for more than two months. In one case they were permanent. In some, after many months, the lesions resolved; in others, complications persisted for three years.^{13/} During the South African epidemic of 1951 several ophthalmologists noted similar changes, but these have not been reported in detail.^{43/}

The relationship between Rift Valley fever and visual sequelae has not been definitely established. Most of the observations reported were on patients without continuous history. Diagnosis of

TABLE V. OCULAR SEQUELAE IN HUMAN CASES OF RIFT VALLEY FEVER

CASE ^a / NUMBER	AGE	OCCUPATION	CONTACT WITH ANIMALS	ONSET OF EYE SYMPTOMS AFTER PYREXIAL ATTACK	FUNDAL FINDINGS	PERSISTENCE OF VISUAL DEFECTS	CF TEST
1	34	Teacher	No	1 week	Large macular mass & small hemorrhage in left eye	>2 months	+
2	34	Research station worker	Yes	1 week	Large macular mass in left eye	>2 months	+
3	29	Research station worker	Yes	3 weeks	Very small macular exudate in left eye	>3 months	+
4	28	Merchant	No	2 weeks	Large paramacular mass in left eye	>1 month	+
5	40	Diamond classifier	No	2-3 days	Large paramacular mass in right eye	Between 6 and 12 weeks	+
6	50	Veterinary surgeon	Yes	Same time	Retinal detachment in right eye	Permanent	+

a. All cases were in males.
Modified from Schrire.^{45/}

Rift Valley fever was made serologically, in most instances weeks or months after the discovery of past illness related by the patient. In three of the six cases reported by Schrire there was, however, a history of contact with animals. Two patients (Cases 2 and 3) were employed at a government research station where a number of cattle were found to have Rift Valley fever. At the same time five other staff members had an influenza-like disease. The third case (Case 6) was a government veterinarian who investigated an outbreak in cattle. This evidence, coupled with the complement-fixation tests, circumstantially linked Rift Valley fever with the reported visual disturbances.

(2) Circulatory

Continuous history of the only reported fatality associated with Rift Valley fever was also inconclusive. Schwentker and Rivers^{39/} reported the history of a pathologist, age 30 years, who contracted the disease in the laboratory. The patient exhibited a typical clinical case of infection and recovery until the sixteenth day after onset, when phlebitis developed in the popliteal vein of the left leg. On days 20 and 26 pulmonary infarcts formed in the right lung. Another infarct formed in the left lung on day 34 and phlebitis developed in the femoral vein of the right leg four days later. The patient died from a large pulmonary embolus on day 45. Autopsy revealed none of the pathological changes associated with Rift Valley fever infections of larger animals. Typically, virus was isolated from the blood on days 1 through 3, but not thereafter. Joubert *et al.*^{48/} referred to one individual who developed coronary thrombosis one week after contracting Rift Valley fever, but stated that the patient was of an age and weight frequently associated with circulatory disorders.

6. Immunity

a. Duration

Antibodies appeared in the sera of infected persons 14 days after onset^{49/} but the duration of active immunity in man following Rift Valley fever infection has not been established.^{50/} Findlay^{51/} reported antibodies in the sera of laboratory workers who had recovered from infection four to five years earlier, but they had had subsequent contact with virus. Sabin and Blumberg^{35/} reported antibodies in the serum of a patient infected and without known contact with the virus 12 years after recovery. Findlay and Howard^{52/} reported on the duration of immunity in the laboratory workers first reported in 1936. The sera of all the workers retained antibodies, which had persisted in the absence of exposure to the virus for periods of 12, 18, and 20 years. Schrire and Gear^{43/} claimed that antibodies persisted for 20 years following infection, and suggested that recovery might be followed by life-long immunity. Brown *et al.*^{50/} questioned the 20-year figure because protocols were not reported, but presented evidence of antibodies persisting for 25 years. Neutralizing

[illegible]

~1758 [102] 102 102

14

antibodies were demonstrated in the serum of one of the authors (T. Dalling) 25 years after recovery from infection. Expressed as the reciprocal of the 50 per cent neutralizing dilution of 250 mouse subcutaneous doses, the serum titer was $1 \times 10^{1.8}$. Dalling had no further contact with the virus after the infection.

b. Immunization

(1) Passive

Findlay^{22/} apparently avoided clinical infection by administering immune human serum during the period of exposure to the virus. He reported the appearance of previously undetected antibodies in his serum without experiencing apparent infection.

(2) Active

Randall^{53/} developed a vaccine for Rift Valley fever, using formalin-inactivated virus which propagated in monkey kidney-cell tissue culture. The vaccine was tested in animals and human volunteers. Neutralizing antibody titers in animals and volunteers were comparable. Antibody titer and immunity were correlated by intracerebral challenge in mice.

Vaccine produced from monkey kidney-cell tissue culture was prepared by inoculating cells in serum-free medium 199 with virus and incubating for 96 to 144 hours at 36°C. Infected whole cultures were homogenized in a Waring blender and clarified by centrifugation. The supernatant blend was passed through a sintered glass filter of medium porosity and the pH was adjusted to 7.0 to 7.2. If the material was of high titer ($10^{8.3}$ to $10^{8.9}$ per ml), it was treated with formalin in a final concentration of 1:1000. Free formalin was neutralized with sodium bisulfite and the vaccine was then dialyzed against Hank's balanced salt solution for 24 hours at 4°C. The vaccine was tested for bacterial sterility and viable virus at several time intervals and then safety-tested in the final procedure.

E. ANIMAL INFECTIONS

1. Natural Infections

Alexander^{8/} and Dickson^{9/} classified Rift Valley fever in sheep and cattle according to clinical signs of the disease. They recognized a peracute form, an acute form, a sub-acute form, and a mild or inapparent form. The peracute form was common in very young lambs. An incubation period of about 12 hours was followed by collapse and death within 36 hours in 95 to 100 per cent of the infected lambs. During the 24 hours preceding death, lambs were listless, disinclined to feed, and sank down soon after being put on their feet. The absence of diarrhea was noted. The acute form was

commonly encountered in lambs and to a lesser extent in adult sheep. Clinical signs appeared suddenly and included a rapid rise in temperature, vomiting, mucopurulent discharge from the nose, rapid pulse, unsteady gait, and abortion in pregnant animals. Death usually followed onset within 24 to 48 hours. The mortality rate was high in lambs and varied from 20 to 30 per cent in adult sheep. In adult sheep and cattle, the subacute form was common. Body temperatures rose to 104° to 106°F and persisted for 24 to 96 hours. There was inappetence and general weakness. Abortion was frequently the only sign in pregnant animals. Milk production decreased rapidly. The mortality rate was low, less than 10 per cent in cattle. The mild or inapparent form also occurred in adult sheep and cattle. The only sign of disease was a mild febrile reaction, and diagnosis could be made only by serological methods. Leucopenia followed the apparent forms of infection.^{22/}

2. Experimental Infections

Experimental infection of lambs with pantropic Rift Valley fever virus (PRVVFV) was followed by a clinical picture similar to natural infections.^{1,22/} Goats showed signs similar to those of sheep. Cattle showed dullness, inappetence and blood-stained nasal discharge.^{55/} Sheep and lambs inoculated with neurotropic Rift Valley fever virus (NRVVFV), other than intracerebrally (IC), exhibited a mild or inapparent form of the disease.^{17,56/} However, when inoculated IC, NRVVFV caused a rapidly fatal encephalitis. Death occurred within 24 hours without signs other than a slightly elevated temperature, but if the animal survived 48 to 72 hours, the elevated temperature was followed by retraction of the head, inability to rise, and convulsive twitching of the limbs.

Mice and rats experimentally infected with NRVVFV and PRVVFV exhibited roughened coats, lethargy, tremors, convulsions, subnormal temperatures in later stages, coma, and usually death in one to three hours after onset of disease and within 36 to 72 hours after inoculation.^{36,57/} Pregnant mice and rats frequently aborted and died. The young were still-born or died shortly after birth.^{17/}

A febrile, nonfatal form of RVF developed in monkeys within 24 to 96 hours after inoculation.^{25/} The febrile response persisted for 24 to 120 hours, followed by leucopenia, but other signs of disease were not apparent.

3. Pathology

Pathology of Rift Valley fever infection in various animals was investigated by Daubney et al.,^{1/} Findlay et al.,^{22,52,56,58/} Marschal,^{59/} Smithburn,^{57/} Kitchen,^{34/} Schulz,^{60/} and Mims^{61/} then reviewed by Weiss in 1957.^{3/} The material presented below was summarized from the observations published by these investigators.

a. Sheep, Cattle, and Goats

The pantropic virus primarily affected the liver, which showed characteristic focal necrosis.^{3/} Liver degeneration in lambs differed somewhat from that in sheep. In lambs, the liver was frequently yellow, rarely enlarged, but always lacking the deep red of the normal liver. It showed necrotic foci approximately one millimeter in diameter in association with subcapsular hemorrhages scattered beneath the capsule. The lesions extended throughout the liver and in peracute cases the normal architecture was sometimes completely lost. The parenchymatous cells underwent hyaline degeneration and lost their affinity for acidophilic stains. With the exception of a few cells near the central vein, the lobule was made up of irregular masses of lightly stained cytoplasm. Leucocytes and histiocytes showing karyorrhexis infiltrated between these masses.

The liver of mature sheep was usually mottled brown and frequently enlarged. Liver cells degenerated and the lesions accumulated polymorphonuclear leucocytes and histiocytes. Lesions were usually focal and not panlobular, as in lambs. Early changes of the liver were observed in a few cells in the central zone of the lobule. Hyaline bodies, indistinguishable from Councilman lesions of yellow fever, were formed by cloudy swelling of the cytoplasm, which was followed by hyaline degeneration. As the lesion progressed, the cells contracted and showed oxychromatic degeneration of the nucleus and development of inclusion bodies. Findlay^{58/} regarded the inclusion bodies as degeneration products. Progressively, the lesions were then invaded by polymorphonuclear leucocytes and histiocytes, which were destroyed and formed a necrotic mass. When the lesion was in the central zone this mass formed an occlusion of the central vein. Generally, liver lesions in cattle and goats were similar to those observed in older sheep.

The spleen in this group of animals usually showed subcapsular petechiae and capillary arborescence, primarily near the free borders. According to Schulz,^{60/} lesions observed during the Union of South Africa epizootic showed, as a rule, tumor splenis and subcapsular hemorrhages. Necrobiotic changes in the pulpa could be seen, along with occasional infiltration of neutrophils.

The kidneys of lambs showed congestion of the cortical and medullary blood vessels, especially near the boundary zone. They showed cloudy swelling, and on occasion the cells of the convoluted tubules lost their ability to retain nuclear stains. Lesions in older sheep most often progressed to tubular degeneration or nephrosis.

The alimentary tract was inflamed in degrees varying from catarrhal to hemorrhagic enteritis. Schulz found large subperitoneal hemorrhages along the entire gastrointestinal tract. In addition to the usual enteritis, he also observed areas of croupous or necrotic enteritis and ulceration.

The wall of the gall bladder showed petechiae; it was often thickened from subserosal and muscular hemorrhage, hyperemia, and edema; it showed extensive desquamation and necrosis of the mucous membrane.

Cyanosis was observed in visible mucous membranes and skin, particularly in the udder, scrotum and axillary regions, the lower part of the extremities, and inside the hind legs. Subcutaneous tissues were edematous and the cutaneous blood vessels were distended. In addition, cattle showed acute catarrhal stomatitis, erosion of the lips, tongue, and cheeks; coronitis, laminitis, exungulation, and, on occasion, marked ascites.^{60/}

In the lungs, congestion of the meningeal vessels and inter-alveolar capillaries was noted. The lungs showed hyperemia, edema and emphysema and subpleural or perivascular hemorrhages. Schulz also observed signs of fibrinous pneumonia in lambs. Degenerative changes were observed in the adrenals. Cortical and medullary hemorrhages and a number of cells containing hyaline inclusions were observed.

The heart showed subpericardial hemorrhages in the region of the coronary grooves and small subendocardial extravasations in the left ventricle. The mesenteric and omental vessels were deeply engorged, and the mesenteric lymph glands were enlarged and moist. At times hemorrhages extended into the cortex of the glands.

The only abnormal change noted in the placenta was invasion of the uterine musculature and decidua by polymorphonuclear leucocytes, many of which were breaking down and undergoing karyorrhexis. Kaschula^{17/} isolated virus from a foetus aborted by an infected ewe and concluded that the virus could pass the placental tissues.

Neurotropic virus, inoculated intracerebrally, produced encephalitis in lambs.^{56/} Focal necrosis, degeneration of ganglion cells and nuclear inclusions in ganglion cells, perivascular infiltration, and infiltration of the meninges with leucocytes were observed.

Pantropic virus produced discrete necrotic foci irregularly distributed throughout the liver lobules of monkeys.^{22/} Foci varied in number, and hyaline degeneration of the cytoplasm was not as well marked as in sheep and goats. Neurotropic virus, inoculated intracerebrally, gave rise to encephalitis similar to that described in sheep.^{56/}

b. Mice

According to Weiss,^{3/} liver lesions were found to be essentially the same in all susceptible species of animals. Findlay,^{22/} Smithburn,^{57/} Kitchen,^{34/} and Mims^{61,62/} described pathology of the disease in mice. The appearance of the liver in infected mice resembled that of lambs; infected

rats showed less extensive damage.^{26/} Smithburn^{57/} and Kitchen^{34/} described encephalitic lesions in mice, similar to those described in lambs and monkeys, when neurotropic virus was inoculated intracerebrally into mice. Fixed neurotropic virus did not produce liver lesions in mice, but virus in its eighty-fifth mouse intracerebral passage in mice produced small foci of degeneration on the livers of mice inoculated intraperitoneally.

Mims^{63/} reported that mice ill with Rift Valley fever had little or no prothrombin in the plasma. Prothrombin levels of less than five per cent of normal and clotting times of 10 to more than 60 minutes were recorded. Mims attributed the hemorrhagic phenomena in Rift Valley fever infections in mice to this deficiency. He pointed out that the deficiency had also been reported in yellow fever infections of rhesus monkeys.

c. Miscellaneous Animals

Findlay^{22/} reported that post-mortem appearances in small rodents such as hamsters, dormice, wood mice, and field moles were very similar to those in lambs and sheep dying of infection with pantropic virus. Marschal^{59/} described histological changes in the livers of hamsters and mice infected with Rift Valley fever. His findings essentially confirmed the observations of Daubney^{1/} and Findlay.^{22/}

Cats showed very small foci of degeneration on the liver. Ferrets presented a pathological picture characterized by extensive development of edematous pulmonary consolidation with a scanty exudate of large mononuclear cells.^{27/}

4. Immunization

a. Passive

Administration of immune serum within 36 hours after exposure to Rift Valley fever virus was shown to protect newly born lambs.^{64/} Practically, this prophylactic method had many disadvantages, and a live or attenuated vaccine that would produce lasting immunity was needed.

b. Active

MacKenzie^{65/} prepared two vaccines, one by inactivation of virus with methylene blue in light and one by inactivation with formalin. A number of investigators studied the attenuation of virus undergoing serial intracerebral passage in mice.^{3/} Kaschula^{17/} found that live vaccine prepared from 86 mouse passages and ten egg passages caused abortion in pregnant animals. Live vaccine prepared from virus with 102 mouse passages and 54 egg passages appeared safe for use in cattle and sheep, although it was slightly less immunogenic than that with fewer passages. According to Weiss,^{3/} the latter vaccine has been used extensively in Africa with encouraging results.

1. The first part of the document is a letter from the President of the United States to the Congress, dated January 1, 1861. It is a formal address, and it is the first of its kind since the signing of the Constitution. The President, James Buchanan, is addressing the Congress, and he is doing so in a very formal and dignified manner. He is discussing the state of the Union, and he is discussing the issues that are facing the country at that time. He is also discussing the role of the President, and he is discussing the responsibilities that he has as the President of the United States. The letter is a very important document, and it is a very interesting one to read. It gives us a glimpse into the mind of the President, and it gives us a glimpse into the state of the country at that time. It is a document that is worth reading, and it is a document that is worth studying.

F. DIAGNOSIS

Confirmed diagnosis of Rift Valley fever has been limited to laboratory methods such as isolation and identification of virus, serological tests, gross pathology, and histopathological studies, especially of the liver.^{3/} Reliance upon symptomatic criteria resulted in delayed recognition of the 1950-1951 outbreak in the Union of South Africa.^{10/} Rift Valley fever was initially confused with diseases such as enterotoxemia and bluetongue in animals and influenza in humans. Similarities to "three-day stiff-sickness" of animals and dengue, yellow fever, and sandfly fever in humans have also been described.^{66/}

G. TREATMENT

Specific treatment of Rift Valley fever has not been reported. Thus, in practice, treatment has been symptomatic.^{3/} Chemotherapeutics tried without success included cortisone and adrenocorticotrophic hormone,^{67/} and prontosil, sulphanilimide, and allied drugs.^{68/} Partial activity against infections in mice treated with chloramphenicol^{69/} and some of the acridines^{70/} has been reported.

III. RIFT VALLEY FEVER VIRUS

A. RECOGNIZED STRAINS

1. Pantropic

RVFV isolated from its natural environment, i.e., an epizootic in African sheep or cattle populations, showed definite affinity for tissues derived from two and possibly three germ layers.^{3/} The virus produced extensive necrosis of liver parenchyma and in this sense was frequently referred to as viscerotropic and hepatotropic. Lesions in tissues of mesodermal origin were of questionable RVFV etiology; the term polytropic was proposed as a more accurate term than pantropic, which was originally applied to viral pathogenesis of all three germ layers.^{34/} The term "polytropic" has not been found in RVF literature other than in the publication in which it was proposed.

2. Neurotropic

A neurotropic variant (NRVFV) of PRVFV was obtained by IC passage in young mice.^{34,57,71,72/} A summary of changes in the virus during 90 passages through mouse brain is given below.

<u>NUMBERED IC PASSAGES IN MICE</u>	<u>CHANGES IN VIRUS AFFINITY FOR MOUSE TISSUE</u>
1-15	No apparent change.
15-20	Gained neurotropism and retained hepatotropism.
20	Hepatotropism lost but easily regained.
33-49	Neurotropism became more stable but could still be lost.
80-90	Neurotropic. Hepatotropism could not be completely regained.

Adapted from Kitchen.^{34/}

The sixty-eighth passage of virus isolated from a human on the first day of illness showed both tropisms; virus isolated on the fourth day exhibited only the hepatic attribute.^{34/} The circulation of eightieth-mouse-passage virus in rhesus monkeys suggested that low-passage neurotropic virus tended to revert to its original pantropic character when injected into monkeys.^{3/}

3. Lunyo Virus

In 1956 Weinbren *et al.*^{32/} isolated two strains of an agent similar to RVFV from mosquitoes caught in the Lunyo Forest on Entebbe peninsula. The virus produced hepatic lesions similar to those produced by PRVSV in mice, but the histopathological picture was different and there were serological and behavioral differences. A summary of similarities and differences between RVFV and Lunyo virus is presented below.

SIMILARITIES	DIFFERENCES
1. Lunyo antiserum neutralized RVFV.	1. RVFV antiserum did not neutralize Lunyo virus.
2. Lunyo virus with strong neurotropic properties readily yielded a viscerotropic strain with IP passage in mice.	2. Lunyo virus was transmitted by <u>Aedes aegypti</u> mosquito, RVFV was not.
3. Lunyo viscerotropic strain showed cross neutralization with PRVSV.	3. Mice infected with neurotropic Lunyo virus became hyperactive, ate any available material including their own appendages, and usually died in convulsions 5 to 12 days after inoculation. Mice infected with NRVSV exhibited inappetence, progressive loss of muscular control, and died within 2 to 3 days after inoculation.
4. Lunyo viscerotropic virus reverted to neurotropic when passaged through a mosquito. PRVSV reacted similarly when passaged through certain rodents.	4. Lunyo virus was less stable in storage (lyophilized in serum) than RVFV.
	5. Lunyo virus failed to yield a hemagglutinin. RVFV readily yielded hemagglutinin.

Adapted from Weinbren *et al.*^{32/}

B. MORPHOLOGY

PRVSV readily passed through Chamberland filters up to L11 grade but rarely passed through filter grade L13.^{1/} By filtration through gradocol membranes, particle size of PRVSV was estimated at 23 to 25 millimicrons.^{49/} Completely different estimates of RVFV particle size were obtained by ultracentrifugation.^{73-75/} PRVSV particle size was reported as 49.7 millimicrons and NRVSV (102 IC mouse passages, 50 egg passages, and 9 IC mouse passages) showed two particle sizes, 30.9 and 51.8 millimicrons.^{73,75/} NRVSV purified by ultracentrifugation remained viable longer than NRVSV in mouse brain preparations.^{76/}

C. PROPAGATION

1. Mice

Mice and rats of several varieties have been used to propagate PRV_{SV}.^{22/} Mims^{36/} used White Swiss mice for propagation and quantitation of RVFV. The virus is 98 to 100 per cent lethal for mice in two to three days.^{22,36/} Mims showed that age of mice had little effect on peak titers in mice 2, 4, 6, 8, and 10 weeks old. The rate of blood titer increase was proportional to the quantity of virus in the inoculum, but the final titer was approximately the same. Peak titers were reached one to two hours before mice sickened and died. Mice inoculated intravenously (IV) or intracerebrally (IC) with 10^2 to 10^3 LD₅₀ per ml* virus produced blood titers as high as 10^{10} LD₅₀ per ml. Subcutaneous (SC) or intraperitoneal (IP) inoculation gave titers in blood approximately one log lower. Titers in brain and liver were usually one to two log₁₀ lower than titers in blood.

2. Lambs

Daubney *et al.*^{1/} used the lamb for isolation and propagation of PRV_{SV}. The virus infection was fatal in two to three days for 90 per cent of the three to seven-day-old lambs. The mortality rate decreased with age to approximately 20 per cent in mature sheep. Exact titers of PRV_{SV} in lambs were not reported, but data presented by Smithburn^{57/} suggested that virus in the bloodstream was plentiful. Two-tenths ml of lamb blood drawn daily from the first through the third days of infection killed all of the mice inoculated intraperitoneally. The lambs were infected with approximately 1×10^4 mouse LD₅₀ of virus given subcutaneously. Neurotropic RVFV was rarely fatal to lambs except when inoculated intracerebrally.^{56/} Smithburn was unable to demonstrate NRV_{SV} in the blood of sheep or lambs during the first ten days following subcutaneous inoculation with approximately 1×10^4 mouse LD₅₀ of NRV_{SV}.^{57/} Availability of lambs was seasonal, however, and their continuous use for propagation of RVFV was impractical. The gestation period in sheep is approximately 150 days;^{77/} therefore, under natural conditions lambs are available only in the spring and late fall.

3. Embryonated Eggs

Saddington^{78/} propagated PRV_{SV} on the chorioallantoic membrane of 9- to 10-day-old embryonated eggs. Embryos were harvested after five days and the membrane, liner, and amniotic fluid were titrated in mice. Titers were not reported, but it was stated that all three materials contained

* LD₅₀ in mice was calculated by the method of Reed and Muench.

virus; the liver was more consistently infectious for mice than membranes or amniotic fluid. Kaschula^{17/} propagated both pantropic and neurotropic strains of the virus in the yolk-sac and on the chorioallantoic membrane of embryonated eggs. Highest titers of virus ($1 \times 10^{5.5}$ mouse LD₅₀) were obtained at 34°C from 8-day-old eggs inoculated in yolk-sac with $1 \times 10^{2.0}$ mouse LD₅₀ per 0.1 ml virus and harvested at death (approximately 48 hours).

4. Tissue Culture

MacKenzie^{79/} and Saddington^{78/} carried PRV_{SV} through 12 subcultures in 9- to 10-day old chick embryo cells suspended in Tyrode's solution. Virus titers of approximately $1 \times 10^{4.5}$ mouse LD₅₀ were obtained after incubation of virus and cell suspension at 37°C for four to five days.^{79/}

Endo^{80/} reported the development of a neurotropic variant by serial passage of PRV_{SV} in a Maitland-type tissue culture of embryonic mouse brain.

In roller tube tissue culture, Takemori *et al.*^{81-83/} demonstrated the cytopathogenic effect of NRV_{SV} and PRV_{SV} on rat sarcoma cells, human embryo, and rat, mouse, and swine fibroblasts. Weiss^{3/} reported cytopathogenesis of NRV_{SV} in lamb kidney cells grown in roller tube cultures. Weiss confirmed the finding of Takemori *et al.* that the cytopathogenic effect appeared within about two days and maximum virus titer (1×10^6 to 1×10^7 mouse LD₅₀) was reached prior to marked cell destruction, which was complete in four to six days.

PRV_{SV} and NRV_{SV} were titrated in tissue culture by plaque formation on rat sarcoma cells,^{81/} Chang's human liver cells,^{84/} and sheep kidney cells.^{85/} Randall^{53/} propagated PRV_{SV} in monkey kidney cell tissue culture for production of vaccine.

5. Adsorption and Multiplication

a. *In vivo*

Mims^{86/} obtained data for PRV_{SV} growth curves by titration in mice. He reported that most of the virus disappeared from the blood within the first hour. Between five and nine hours, depending upon inoculum size, there was an exponential rise in titer. Comparable final yields of virus (1×10^8 MICLD₅₀ per 0.03 ml) were obtained for different-sized inocula at different times, as shown below.

INOCULUM, MICLD ₅₀ PER 0.03 ml	APPROXIMATE TIME, HOURS
$1 \times 10^{0.7}$	50
$1 \times 10^{1.5}$	48
$1 \times 10^{4.5}$	30
$1 \times 10^{7.3}$	18

Adapted from Mims.^{86/}

...the first was not completely ...
or
... ..
... ..
... ..
... ..

... ..

... ..
... ..
... ..
... ..

... ..

... ..
... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..

Two-step growth curves developed when inocula of 1×10^6 to $1 \times 10^{7.5}$ MICLD₅₀ per 0.03 ml were used. The curve was characterized by an exponential rise at four to five hours, when a one-hour lag appeared that was followed by a second exponential rise. One-step growth curves resulted from increasing the inoculum size to $1 \times 10^{8.3}$ to $1 \times 10^{8.5}$ MICLD₅₀ per 0.03 ml. Titers increased exponentially between four or five hours and seven or eight hours without evidence of lag. Peak titers of $1 \times 10^{8.5}$ to $1 \times 10^{9.0}$ MICLD₅₀ were reached in seven or eight hours.

Takemori *et al.*^{82/} constructed growth curves for both neurotropic and pantropic RVFV from data obtained by titration of ascitic fluid from infected mice that had been inoculated with ascites hepatoma cells eight days before virus inoculation. Matumoto *et al.*^{87/} confirmed the findings with neurotropic RVFV.

b. In vitro

Plowright and Ferris^{85/} reported growth curve studies of RVFV using sheep kidney monolayers in culture tubes. They reported peak titers of $1 \times 10^{6-6.5}$ MIPLD₅₀ in about 36 hours. Takemori *et al.*^{83/} using ascites hepatoma cells of mice in roller tube tissue culture, made quantitative measurements of NRVFV and PRVFV. They found little difference in the in vitro growth of the two virus strains. Peak titers of 1×10^6 MICLD₅₀ were attained after two to three days.

Iwasa^{84/} employed plaque formation on monolayers of Chang's human liver cells (CHL) to investigate the latent period and the appearance of intracellular antigen of RVFV. He stated that the period of intracellular multiplication was 2.5 hours and the latent period extended to 6 hours. Using the indirect fluorescent antibody technique, he detected intracellular complement-fixing antigen at 5 hours.

c. Interference

Findlay and Howard^{52/} investigated interference between neurotropic and hepatotropic virus and between ether-inactivated neurotropic virus and hepatotropic virus. Exaltation between Senger virus and neurotropic virus was also studied. They claimed correlation between the development of neurotropism and increasing interference between neurotropic and hepatotropic strains. Because of difficulties encountered in obtaining completely inactivated virus by ether treatment, these results were not clear-cut, but the authors concluded that ether-inactivated virus interfered with living virus. It was also concluded that Senger virus, a member of the encephalomyocarditis group of viruses, was exalted by neurotropic virus injected intraperitoneally.

Naude and Polson^{88/} investigated interference between active RVFV and ultraviolet-irradiated RVFV. They found that irradiated virus interfered with infectivity and that optimal doses of irradiated virus elicited immunity in mice. Joya^{89/} investigated the antigenic power of RVFV inactivated by UV irradiation.

Scott and Whitcomb^{90/} investigated interference of RVFV by rinderpest virus (RV) in hamsters and mice. Hamsters that were infected with RV two to seven days earlier were not affected by exposure to RVFV. The survivors were found to be susceptible to RVFV when re-exposed. Exposure to RV 8, 9, and 10 days before exposure to RVFV resulted in 50 per cent deaths from delayed RVFV infection. Survivors were immune upon re-exposure to RVFV. Periods of 1, 12, 13, and 14 days pre-exposure of hamsters to RV produced no evidence of interference on inoculation with RVFV. Similar evidence of interference did not occur in mice. The only significant feature of the mouse experiments was a delay in peak death time. Similarity of Rift Valley fever to dengue and yellow fever induced Findlay^{22/} to investigate interference among these viruses. He reported no interference between RVFV and dengue virus. However, Findlay and MacCallum^{91/} reported that IP inoculation of mice with a mixture of PRFV and neurotropic yellow fever virus (NYFV) completely protected a few mice and delayed the death time of the others. This protection was not seen if PRFV was given 24 hours prior to inoculation with NYFV. Similarly, it was shown that NYFV protects against pantropic yellow fever virus (PYFV) but PYFV did not protect against NYFV.

d. Incomplete Virus

Isaacs^{92/} preferred limited use of the term incomplete virus and restricted its application to focusing attention on the differences in the infectivity of virus produced from serial passage of low-dilution and high-dilution material. He criticized use of the term to imply the production of an abnormal end-product of virus multiplication or the production of developmentally incomplete virus. Mims^{93/} ascribed the low infectivity of material produced by serial passage of large inocula of PRFV to the production of incomplete noninfective virus. He found that mouse serum with a PRFV titer of 1×10^9 MIVLD₅₀ per 0.03 ml, passaged undiluted in mice, dropped as low as $1 \times 10^{1.9}$ MIVLD₅₀ per 0.1 ml at the fifth passage. The hemagglutinin titer of this material was not reduced in proportion to the infectivity titer, but remained relatively constant after the first passage. Material diluted to 10^{-8} produced high infectivity titers and proportionately high hemagglutinin titers.

D. STABILITY

1. Acid Tolerance

Findlay^{22/} reported that PRFV in blood adjusted to pH 6.9 to 7.3 with phosphate buffer retained viability for 20 hours at 37°C but was

1800-1801

SECRET

inactivated at room temperature. At pH 6.6 at room temperature, some virus remained viable; however, at pH 8 the virus was inactivated both at room temperature and at 37°C. In contrast, Kaschula^{17/} reported that maximum viability of the virus was sustained at approximately pH 8. Mims^{36/} tested virus stability at various pH values from 6.0 to 8.0 at 37.5°C in Sorensen buffer, Sorensen-buffered saline, bovine albumin, and urea concentrations of 0.3 and 3.0 per cent. He found that the virus was equally stable in all the diluents tested as long as the pH was between 7 and 8. Below pH 6.0 the virus was rapidly inactivated. Mims concluded that the absence of virus from the urine of infected mice could probably be attributed to the pH of mouse urine, normally between pH 4 and 6 and rarely as high as pH 7.

2. Heat

PRVSV in phosphate-buffered blood (pH 7.2) retained virulence for 20 minutes at 56°C but not for 40 minutes, according to Findlay.^{22/} Mims^{94/} found that undiluted mouse serum with a PRVSV titer of 1×10^9 MICLD₅₀ per 0.03 ml lost only one to two logs of titer when heated to 56°C for one hour. Surprisingly, the titer did not drop below 1×10^5 MICLD₅₀ per 0.03 ml when the sample was heated to 56°C for three hours. In contrast, the hemagglutinin of PRVSV was completely destroyed in one hour at 56°C.

3. Chemicals

Daubney et al.^{1/} recovered viable PRVSV from blood plasma after salting out virus and protein fractions with ammonium sulphate buffered at pH 7.4. Andrewes and Horstmann^{95/} treated PRVSV with ethyl ether for 18 to 24 hours at 4°C. They added ether to a mouse liver suspension with a titer of 1×10^4 MIPLD₅₀ per ml and found that the titer was reduced 100-fold. Mims^{94/} found that RVSV extracted from infective mouse serum with acetone and ethyl ether lost its infectivity but retained antigenicity and hemagglutinin.

4. Storage

a. Cold

According to Daubney et al.^{1/} PRVSV in citrated blood may be maintained for one week at room temperature without loss of infectivity. With oxalate-carbol-glycerine (O.C.G.) added, infective blood stored under refrigeration at 5°C (with considerable variation) retained virulence for 54 days and remained viable with reduced virulence for 147 days. Findlay^{22/} reported that infective blood with O.C.G. remained viable for eight months under refrigeration at 4°C, but with reduced virulence. Defibrinated blood with 0.5 per cent phenol remained virulent for six months at 4°C. PRVSV in undiluted mouse serum remained viable for eight months and showed relatively no loss in titer for 30 days at -20°C.^{36/} Luny virus, stored under the same stated conditions, lost 1.2 logs of virus in 28 days.^{32/}



Kaschula^{17/} confirmed Endo's finding^{96/} that RVFV was most stable in whole egg. This stabilizing action was later attributed to egg yolk. Kaschula concluded that the virus was most stable in undiluted egg embryo at -70°C . Instability of RVFV in mouse brain suspension^{97/} was attributed to the action of virus-inactivating enzymes, the activity of which was retarded by 0.01 M KCN, condensation products of gallic acid and formalin, and phosphorylation products of resorcinol.^{3/}

b. Lyophilization

PRVFV in mouse blood dried in vacuo over P₂O₅ and maintained at 4°C remained active for at least six weeks.^{22/} Dried from the frozen state in vacuo over H₂SO₄ and maintained at 4°C , the virus in blood diluted 1:10 with normal serum remained viable for eight months.^{38/} Lyophilized NRVFV in mouse brain lost titer when stored at 37°C for one week or at 4°C for three months. It was found that the presence of buffering electrolytes was harmful to the stability of lyophilized preparations. Stability was improved by the addition of five per cent sucrose, one per cent mixture of 21 amino acids, and five per cent peptone or a saturated solution of lactalbumen hydrolyzate.^{3/} A lyophilized preparation of Lunyo virus in mouse brain with 100 per cent normal serum as the desiccating medium dropped from a titer of $1 \times 10^{6.8}$ MICLD₅₀ to a titer of $1 \times 10^{4.5}$ MICLD₅₀ after storage for six weeks. A similar preparation employing beef-peptone-albumin as the desiccating medium lost 4.2 logs of potency after 18 days' storage at -20°C .

E. IDENTIFICATION PROCEDURE

1. Isolation

Rift Valley fever virus has been successfully isolated by injection of suspected material into highly susceptible animals such as lambs or sheep,^{1/} mice,^{72/} and ferrets.^{27/} Isolation has also been made by inoculation of infected material into the yolk-sac or on the chorioallantoic membrane of eight-day embryonated eggs.^{17/} Whole blood,^{1/} serum,^{22/} throat and nasal washings,^{27/} homogenates of insects,^{5/} egg embryo,^{17/} and vital organs such as brain, liver and spleen from suspected sources have been successfully used as inocula.^{22/} Virus has been concentrated from suspension by precipitation and filtration.^{1/} Whole blood and serum, aseptically drawn from disease-free animals, have been widely used to obtain bacteria-free suspensions of virus. The method requires knowledge of when and where virus is found in various animals. It has been shown^{98/} that the titer of pantropic virus in various organs was related to concentration of blood in the organs. The periods during which the more important host species have been found to have virus in the blood are compiled in Table VI.

TABLE VI. CIRCULATION OF PRV FV IN BLOOD OF VARIOUS HOSTS

HOST	DAYS OF PEAK TITER	DAYS AFTER INOCULATION VIRUS PERSISTS IN BLOOD	LITERATURE CITED
Sheep	3-5	6-7	22
Monkey	4-6	13	22
Rat	3-5	15	22
Human	1-3	9	22
Mice	1-2	2-4	36
Lambs	2-3	2-3	36
Ferrets	2-4	2-4	27

2. Identification

a. Cross-Immunity

Thus far, positive identification of Rift Valley fever virus has been accomplished by the adaptation of accepted serological techniques. The earliest test used for the identification of the virus was introduced by Daubney *et al.*,^{1/} who performed cross-immunity tests on lambs. Seasonal availability, manipulation, and expense incurred in the extensive use of lambs for routine work were serious limitations of this method. Findlay^{99,100/} found no cross-immunity between Rift Valley fever and louping ill or dengue in monkeys.

b. Serum Neutralization

Findlay^{22/} discovered the high susceptibility of mice. He inoculated mice with dilutions of infected blood neutralized with a constant amount of antiserum. Modifications of this test have been applied to differentiation of neurotropic and pantropic strains by intradermal, intracerebral, subcutaneous, and intraperitoneal inoculation.^{34/} Findlay^{51/} also showed that the neutralized virus could be reactivated by simple dilution or nasal instillation. Inactivation of nonspecific inhibitors by heating at 56°C for 30 minutes has been recommended.^{17/}

c. Complement Fixation

The complement-fixation test was adapted to the identification of Rift Valley fever virus and its homologous antisera by Broom and Findlay in 1932.^{101/} They showed that fixation was in direct proportion to severity

STATE OF NEW YORK

NAME	RESIDENCE	DATE
John Doe	New York	1890
Jane Smith	New York	1891
Robert Brown	New York	1892
William Jones	New York	1893
Elizabeth White	New York	1894
Thomas Black	New York	1895
Charles Green	New York	1896
Harriet Lee	New York	1897
James Hall	New York	1898
Anna King	New York	1899
George Baker	New York	1900

1890

1891

1892

1893

1894

1895

1896

1897

1898

1899

1900

of clinical disease. Exact correlation between the serum-neutralization test and the complement-fixation test was reported by Gear *et al.*^{12,13/} It was reported that complement-fixing antibodies were not always detectable in the sera of humans and animals inoculated with the neurotropic strain, but the report was not supported by experimental evidence. Iwasa^{84/} studied the production of complement-fixing antigen in tissue culture (CHL cells), using the fluorescent antibody technique.

d. Hemagglutination Inhibition

Mims and Mason^{94/} adapted the hemagglutination inhibition test (HAI) to the identification of Rift Valley fever virus. They showed a specificity of the test for the virus and no serological overlap with Semliki forest (Group A) or yellow fever (Group B) viruses. Optimal hemagglutination occurred at pH 6.5 and 25°C. Acetone-ether-treated and untreated virus were adsorbed to red cells. The virus was not eluted, nor could it be washed off. The untreated virus preparation was a more powerful hemagglutinin than the acetone-ether-treated preparation, which was noninfective. Chick red blood cells were routinely used for the test.

e. Agar Diffusion

Although substantiating evidence was not presented, Weiss^{3/} reported demonstration of a precipitin reaction with neurotropic virus and its antibody in agar plates.

F. CLASSIFICATION

Casals^{102/} grouped arthropod-borne viruses into three groups (A, B, and C) on the basis of serological interactions. A fourth group (unclassified) listed viruses that were not placed in the A, B, or C groups because they failed to show serologic relationship to any of the other arthropod-borne viruses. Rift Valley fever virus was placed in the unclassified group. Polson and Linder^{74,75/} grouped eight animal viruses into three groups on the basis of ultracentrifuge sedimentation rates of 70, 170, and 450 Svedbergs. Rift Valley fever virus fell into the 450-Svedberg group, along with neurotropic horse-sickness virus. Bauer and Bradley^{103/} classified neurotropic viruses according to xanthine oxidase activity in infected mouse brain. They placed neurotropic Rift Valley fever virus in group two of their five groups. Bunyamwera, encephalomyelitis, neurovaccinia, pseudolymphocytic choriomeningitis, Fantz and California viruses were also placed in this group.

The first of these is the fact that the
 Government has not yet decided whether
 it will accept the offer of the
 United States to purchase the
 Alaska Pipeline. This is a
 very important decision, and
 it will have a great effect
 on the future of the
 country.

4. For the following:

It is the intention of the Board of Directors of this bank to
 issue a new series of shares of common stock, the proceeds of which
 will be used for the purpose of increasing the capital of the bank
 and for the purpose of providing for the redemption of the shares
 of common stock which are now outstanding. The Board of Directors
 has authorized the issuance of such shares and has authorized the
 officers of the bank to execute all necessary documents in connection
 with the issuance of such shares.

[illegible]

142

IV. SUMMARY

Daubney et al.^{1/} and Weiss^{3/} suggested that RVF was present in Central Africa for years before 1930. Montgomery (1912) and Stordy (1913) described a similar disease in the Rift Valley, but in neither instance was the etiological agent determined.

Findlay and co-workers described the pathology and histopathology of the disease and adapted the serum-neutralization test and the complement-fixation test to detection of RVFV antibodies in serum in 1932.^{22/} In 1934 Kitchen made a detailed report on the disease in man.^{38/} In 1936 MacKenzie and Findlay developed a neurotropic strain (NRVFFV) of the virus by serial passage in mouse brain.^{72/} In the same year the complement-fixation test was used to outline the geographical area of the disease in Africa by extensive testing of sera from African natives.^{6/}

Smithburn et al.^{5/} established arthropod transmission of the virus in 1948 by isolating it from mosquitoes caught in the uninhabited Semliki forest in Western Uganda. Subsequently, in 1949, they succeeded in passing the virus in experimental animals by mosquito bite.^{28/} In 1950-1951 the first known epizootic of the disease occurred in South Africa.^{104/}

Takemori et al.^{81/} reported plaque formation by RVFV in monolayer tissue culture in 1955. In 1956, Weinbren et al.^{32/} isolated the Lunyo virus, which produces cannibalism in mice. RVF epizootics reappeared in local areas of Africa in 1953, 1955, and 1956.^{3/} At this writing, the disease has not been reported under natural circumstances outside Africa. Laboratory infections^{2/} have been reported in Africa, England, Japan, and the United States.

LITERATURE CITED

1. Daubney, R.; Hudson, J.R.; and Garnham, P.C.: "Enzootic Hepatitis or Rift Valley Fever; An Undescribed Virus Disease of Sheep, Cattle and Man from East Africa," J Pathol Bacteriol, 34:4:545-579, 1931.
2. Todd, F.A.; Battles, R.; Beaudette, F.R.; Carlson, E.E.; DeTray, D.E.; Hendershott, R.A.; Mitchell, C.A.; Schroeder, C.R.; Kuttler, K.A.; Shahan, M.S.; Traum, J.; and Winter, A.: "Rift Valley Fever in Foreign Animal Diseases: Their Prevention, Diagnosis and Control," Report of Committee on U.S. Livestock Sanitary Assn, 149-158, 1954.
3. Weiss, K.E.: "Rift Valley Fever - A Review," Bull Epiz Dis Africa, 5:4:431-458, 1957.
4. Kaschula, V.R.: "Comparison of Important Exotic Diseases with Common Indigenous Diseases of the U.S.A.," J S African Vet Med Assoc, 27:3: 201-208, 1956.
5. Smithburn, K.C.; Haddow, A.J.; and Gillett, J.D.: "Rift Valley Fever. Isolation of the Virus from Wild Mosquitoes," Brit J Exptl Pathol, 29:2:107-121, 1948.
6. Findlay, G.M.; Stefanopoulo, G.M.; and MacCallum, F.O.: "Presence d'Anticorps Contre la Fievre de la Vallee du Rift Dans le Sang des Africains," Bull Soc Pathol Exotique, 29:986-996, 1936.
7. Daubney, R., and Hudson, J.R.: "Rift Valley Fever," E African Med J, 10:2-19, 1933.
8. Alexander, R.A.: "Rift Valley Fever in the Union," J S African Vet Med Assoc, 22:3:105-109, 1951.
9. Dickson, J.L.: "Rift Valley Fever in the Union," J S African Vet Med Assoc, 22:3:110-111, 1951.
10. Schulz, K.: "Rift Valley Fever in South Africa: Appreciation of Public Health Significance with Prognostication of Possible Future Epidemics and the Possibility of Vector Control," Special Rpt 5/51, Union Dept of Health Plague Research Laboratory, 1951.
11. Gear, J.: "Studies on Rift Valley Fever," Proc Transvaal Soc Pathol, October 1953.
12. Gear, J.; deMeillon, B.; Measrock, V.; Harwin, R.; and Davis, D.H.S.: "Rift Valley Fever in South Africa. II. The Occurrence of Human Cases in the Orange Free State, the North-Western Cape Province, The Western and Southern Transvaal. B. Field and Laboratory Investigations," S African Med J, 25:49:908-912, 1951.

1. The above is a list of the names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900.

2. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

3. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

4. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

5. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

6. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

7. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

8. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

9. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

10. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

11. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

12. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

13. The names of the persons who have been appointed to the various committees of the Board of Directors of the United States National Bank of Commerce, New York City, for the year ending December 31, 1900, are as follows:

13. Gear, J.; deMeillon, B.; LeRoux, A.F.; Kofsky, R.; Rose-Innes, R.; Steyn, J.J.; Oliff, W.D.; and Schulz, K.H.: "Rift Valley Fever in South Africa. A Study of the 1953 Outbreak in Orange Free State with Special Reference to Vectors and Possible Reservoir Hosts," S African Med J, 29:22:514-518, 1955.
14. Kokernot, R.H.; Heymann, C.S.; Muspratt, J.; and Wolstenholme, B.: "Studies on Arthropod-Borne Viruses of Tongaland. V. Isolation of Bunyamwera and Rift Valley Fever Viruses from Mosquitoes," S African J Med Sci, 22:71-80, 1957.
15. Shone, D.K.: "Rift Valley Fever in Southern Rhodesia," Central African J Med, 4:7:284-286, 1958.
16. Van der Linde, N.T.: "A Recent Epidemic of Rift Valley Fever in the Orange Free State," J S African Vet Med Assoc, 24:3:145-148, 1953.
17. Kaschula, V.R.: "The Propagation and Modification of Strains of Rift Valley Fever Viruses in Embryonated Eggs and Their Use as Immunizing Agents for Domestic Ruminants," Thesis, University of Pretoria, 1953.
18. Kokernot, R.H.; Smithburn, K.C.; and Weinbren, M.P.: "Neutralizing Antibodies to Arthropod-Borne Viruses in Human Beings and Animals in the Union of South Africa," J Immunol, 77:313-323, 1956.
19. Kokernot, R.H.; Smithburn, K.C.; Weinbren, M.P.; and deMeillon, B.: "Studies on Arthropod-Borne Viruses of Tongaland. VI. Isolation of Pongola Virus from Aedes (Banksinella) circumluteolus Theo," S African J Med Sci, 22:81-92, 1957.
20. Kokernot, R.H.; Smithburn, K.C.; Muspratt, J.; and Hodgson, B.: "Studies on Arthropod-Borne Viruses of Tongaland. VIII. Spondweni Virus, an Agent Previously Unknown, Isolated from Taeniorhynchus (Mansonioides) uniformis Theo," S African J Med Sci, 22:103-112, 1957.
21. Paterson, H.E.; Kokernot, R.H.; and Davis, D.H.S.: "Studies on Arthropod-Borne Viruses of Tongaland. IV. The Birds of Tongaland and Their Possible Role in Virus Disease," S African J Med Sci, 22:63-69, 1957.
22. Findlay, G.M.: "Rift Valley Fever or Enzootic Hepatitis," Trans Roy Soc Trop Med Hyg, 25:4:229-265, 1932.
23. Findlay, G.M., and Daubney, R.: "The Virus of Rift Valley Fever or Enzootic Hepatitis," Lancet, 221:5651:1350-1351, 1931.
24. MacKenzie, R.D.; Findlay, G.M.; and Stern, R.O.: "Studies on Neurotropic Rift Valley Fever Virus. The Susceptibility of Rodents," Brit J Exptl Pathol, 17:352-361, 1936.

1. The first part of the report deals with the general situation of the country and the progress of the work during the year.
2. The second part deals with the results of the work done during the year.
3. The third part deals with the financial statement of the year.
4. The fourth part deals with the general remarks of the committee.
5. The fifth part deals with the conclusions of the committee.
6. The sixth part deals with the recommendations of the committee.
7. The seventh part deals with the general remarks of the committee.
8. The eighth part deals with the conclusions of the committee.
9. The ninth part deals with the recommendations of the committee.
10. The tenth part deals with the general remarks of the committee.
11. The eleventh part deals with the conclusions of the committee.
12. The twelfth part deals with the recommendations of the committee.
13. The thirteenth part deals with the general remarks of the committee.
14. The fourteenth part deals with the conclusions of the committee.
15. The fifteenth part deals with the recommendations of the committee.
16. The sixteenth part deals with the general remarks of the committee.
17. The seventeenth part deals with the conclusions of the committee.
18. The eighteenth part deals with the recommendations of the committee.
19. The nineteenth part deals with the general remarks of the committee.
20. The twentieth part deals with the conclusions of the committee.
21. The twenty-first part deals with the recommendations of the committee.
22. The twenty-second part deals with the general remarks of the committee.
23. The twenty-third part deals with the conclusions of the committee.
24. The twenty-fourth part deals with the recommendations of the committee.
25. The twenty-fifth part deals with the general remarks of the committee.
26. The twenty-sixth part deals with the conclusions of the committee.
27. The twenty-seventh part deals with the recommendations of the committee.
28. The twenty-eighth part deals with the general remarks of the committee.
29. The twenty-ninth part deals with the conclusions of the committee.
30. The thirtieth part deals with the recommendations of the committee.
31. The thirty-first part deals with the general remarks of the committee.
32. The thirty-second part deals with the conclusions of the committee.
33. The thirty-third part deals with the recommendations of the committee.
34. The thirty-fourth part deals with the general remarks of the committee.
35. The thirty-fifth part deals with the conclusions of the committee.
36. The thirty-sixth part deals with the recommendations of the committee.
37. The thirty-seventh part deals with the general remarks of the committee.
38. The thirty-eighth part deals with the conclusions of the committee.
39. The thirty-ninth part deals with the recommendations of the committee.
40. The fortieth part deals with the general remarks of the committee.
41. The forty-first part deals with the conclusions of the committee.
42. The forty-second part deals with the recommendations of the committee.
43. The forty-third part deals with the general remarks of the committee.
44. The forty-fourth part deals with the conclusions of the committee.
45. The forty-fifth part deals with the recommendations of the committee.
46. The forty-sixth part deals with the general remarks of the committee.
47. The forty-seventh part deals with the conclusions of the committee.
48. The forty-eighth part deals with the recommendations of the committee.
49. The forty-ninth part deals with the general remarks of the committee.
50. The fiftieth part deals with the conclusions of the committee.

25. Findlay, G.M.: "The Infectivity of Rift Valley Fever for Monkeys," Trans Roy Soc Trop Med Hyg, 26:2:161-168, 1932.
26. Findlay, G.M., and Howard, E.M.: "The Susceptibility of Rats to Rift Valley Fever in Relation to Age," Ann Trop Med Parasitol, 46: 1:35-37, 1952.
27. Francis, T., and Magill, T.P.: "Rift Valley Fever: A Report of Three Cases of Laboratory Infection and Experimental Transmission of the Disease to Ferrets," J Exptl Med, 62:3:433-448, 1935.
28. Smithburn, K.C.; Haddow, A.J.; and Lumsden, W.H.R.: "Rift Valley Fever; Transmission of the Virus by Mosquitoes," Brit J Exptl Pathol, 30:1:35-47, 1949.
29. Dick, G.W.A.: "Epidemiological Notes on Some Viruses Isolated in Uganda (Yellow Fever, Rift Valley Fever, Bwamba Fever, West Nile, Mengo, Semliki Forest, Bunyamwera, Ntaya, Uganda S, and Zika Viruses)," Trans Roy Soc Trop Med Hyg, 47:13-43, 1953.
30. Weinbren, M.P.; Heymann, C.S.; Kokernot, R.H.; and Paterson, H.E.: "Studies on Arthropod-Borne Viruses of Tongaland. VII. Simbu Virus, a Hitherto Unknown Agent Isolated from Aedes (Banksinella) circum-luteolus," S African J Med Sci, 22:93-102, 1957.
31. Gillett, J.D., and Mims, C.A.C.: "Rift Valley Fever," E African Virus Res Inst, Report No. 5, 4-7, 1955.
32. Weinbren, M.P.; Williams, M.C.; and Haddow, A.J.: "A Variant of Rift Valley Fever," S African Med J, 31:38:951-957, 1957.
33. Steyn, J.H., and Schulz, K.H.: "Aedes (Ochlerotatus) caballus Theo, the South African Vector of Rift Valley Fever," S African Med J, 29:48:1114-1120, 1955.
34. Kitchen, S.F.: "The Development of Neurotropism in Rift Valley Fever Virus," Ann Trop Med Parasitol, 44:2:132-145, 1950.
35. Sabin, A.B., and Blumberg, R.W.: "Human Infection with Rift Valley Fever Virus and Immunity Twelve Years after Single Attack," Proc Soc Exptl Biol Med, 64:4:385-389, 1947.
36. Mims, C.A.C.: "Rift Valley Fever Virus in Mice. I. General Features of the Infection," Brit J Exptl Pathol, 37:2:99-109, 1956.
37. Weinbren, M.P., and Mason, P.J.: "Rift Valley Fever in a Wild Field Rat (Arvicanthis abyssinicus): A Possible Natural Host," S African Med J, 31:18:427-430, 1957.

23. The first of these is the fact that the
the first of these is the fact that the

24. The second of these is the fact that the
the second of these is the fact that the

25. The third of these is the fact that the
the third of these is the fact that the

26. The fourth of these is the fact that the
the fourth of these is the fact that the

27. The fifth of these is the fact that the
the fifth of these is the fact that the

28. The sixth of these is the fact that the
the sixth of these is the fact that the

29. The seventh of these is the fact that the
the seventh of these is the fact that the

30. The eighth of these is the fact that the
the eighth of these is the fact that the

31. The ninth of these is the fact that the
the ninth of these is the fact that the

38. Kitchen, S.F.: "Laboratory Infections with the Virus of Rift Valley Fever," Am J Trop Med, 14:6:547-564, 1934.
39. Schwentker, F.F., and Rivers, T.M.: "Rift Valley Fever in Man. Report of Fatal Laboratory Infection Complicated by Thrombophlebitis," J Exptl Med, 59:3:305-313, 1934.
40. Smithburn, K.C.; Mahaffy, A.F.; Haddow, A.J.; Kitchen, S.F.; and Smith, J.F.: "Rift Valley Fever; Accidental Infections Among Laboratory Workers," J Immunol, 62:2:213-227, 1949.
41. Matumoto, M.; Iwasa, S.; and Endo, M.: "Complement Fixation Reaction of Rift Valley Fever," Japanese J Exptl Med, 20:501-508, 1950.
42. Mundel, B., and Gear, J.: "Rift Valley Fever. I. The Occurrence of Human Cases in Johannesburg," S African Med J, 25:44:797-800, 1951.
43. Schrire, L., and Gear, J.: "The Fevers of Africa. III. Rift Valley Fever," Central African J Med, 2:6:237-240, 1956.
44. Freed, I.: "Rift Valley Fever in Man Complicated by Retinal Changes and Loss of Vision," S African Med J, 25:50:930-932, 1951.
45. Schrire, L.: "Macular Changes in Rift Valley Fever," S African Med J, 25:50:926-930, 1951.
46. Stern, L.: "Rift Valley Fever in Rhodesia: Report of a Case in a Laboratory Worker," Central African J Med, 4:7:281-283, 1958.
47. Mims, C.A.C.: "Rift Valley Fever: Infections in Laboratory Staff," E African Virus Res Inst, Report No. 5:6-7, 1955.
48. Joubert, J.D.S.; Ferguson, A.L.; and Gear J.: "Rift Valley Fever in South Africa. II. The Occurrence of Human Cases in the Orange Free State, the North-Western Cape Province, the Western and Southern Transvaal. A. Epidemiological and Clinical Findings," S African Med J, 25:48:890-891, 1951.
49. Broom, J.C., and Findlay, G.M.: "The Filtration of Rift Valley Fever Virus Through Graded Collodion Membranes," Brit J Exptl Pathol, 14:3:179-181, 1933.
50. Brown, R.D.; Scott, G.R.; and Dalling, T.: "Persistence of Antibodies to Rift Valley Fever in Man," Lancet, 273:6990-6991:345, 17 August 1957.
51. Findlay, G.M.: "The Mechanism of Immunity in Rift Valley Fever," Brit J Exptl Pathol, 17:89-104, 1936.

1. [Illegible text]
2. [Illegible text]
3. [Illegible text]
4. [Illegible text]
5. [Illegible text]
6. [Illegible text]
7. [Illegible text]
8. [Illegible text]
9. [Illegible text]
10. [Illegible text]
11. [Illegible text]
12. [Illegible text]
13. [Illegible text]
14. [Illegible text]
15. [Illegible text]
16. [Illegible text]
17. [Illegible text]
18. [Illegible text]
19. [Illegible text]
20. [Illegible text]
21. [Illegible text]
22. [Illegible text]
23. [Illegible text]
24. [Illegible text]
25. [Illegible text]
26. [Illegible text]
27. [Illegible text]
28. [Illegible text]
29. [Illegible text]
30. [Illegible text]
31. [Illegible text]
32. [Illegible text]
33. [Illegible text]
34. [Illegible text]
35. [Illegible text]
36. [Illegible text]
37. [Illegible text]
38. [Illegible text]
39. [Illegible text]
40. [Illegible text]
41. [Illegible text]
42. [Illegible text]
43. [Illegible text]
44. [Illegible text]
45. [Illegible text]
46. [Illegible text]
47. [Illegible text]
48. [Illegible text]
49. [Illegible text]
50. [Illegible text]
51. [Illegible text]
52. [Illegible text]
53. [Illegible text]
54. [Illegible text]
55. [Illegible text]
56. [Illegible text]
57. [Illegible text]
58. [Illegible text]
59. [Illegible text]
60. [Illegible text]
61. [Illegible text]
62. [Illegible text]
63. [Illegible text]
64. [Illegible text]
65. [Illegible text]
66. [Illegible text]
67. [Illegible text]
68. [Illegible text]
69. [Illegible text]
70. [Illegible text]
71. [Illegible text]
72. [Illegible text]
73. [Illegible text]
74. [Illegible text]
75. [Illegible text]
76. [Illegible text]
77. [Illegible text]
78. [Illegible text]
79. [Illegible text]
80. [Illegible text]
81. [Illegible text]
82. [Illegible text]
83. [Illegible text]
84. [Illegible text]
85. [Illegible text]
86. [Illegible text]
87. [Illegible text]
88. [Illegible text]
89. [Illegible text]
90. [Illegible text]
91. [Illegible text]
92. [Illegible text]
93. [Illegible text]
94. [Illegible text]
95. [Illegible text]
96. [Illegible text]
97. [Illegible text]
98. [Illegible text]
99. [Illegible text]
100. [Illegible text]

52. Findlay, G.M., and Howard, E.M.: "Notes on Rift Valley Fever," Arch ges Virusforsch, 4:4:411-423, 1948-1952.
53. Randall, R.; Gibbs, C.J.; Aulisio, C.G.; and Binns, L.M.: "Development of a Formalinized Rift Valley Fever Vaccine," Fed Proc, 19:219, 1960.
54. Daubney, R.: "Newer Researches Regarding Tropical and Subtropical Diseases," Report of 13th Int Vet Congress, 2:752-764, 1938.
55. Henning, M.W.: "Rift Valley Fever," J S African Vet Med Assoc, 23:2:65-73, 1952.
56. Findlay, G.M.; MacKenzie, R.D.; and Stern, R.O.: "Studies on Neurotropic Rift Valley Fever Virus: The Susceptibility of Sheep and Monkeys," Brit J Exptl Pathol, 17:6:431-441, 1936.
57. Smithburn, K.C.: "Rift Valley Fever; the Neurotropic Adaptation of the Virus and the Experimental Use of this Modified Virus as a Vaccine," Brit J Exptl Pathol, 30:1:1-16, 1949.
58. Findlay, G.M.: "Cytological Changes in the Liver in Rift Valley Fever, with Special Reference to the Nuclear Inclusions," Brit J Exptl Pathol, 14:4:207-219, 1933.
59. Marschal, F.: "Die Histologischen Leberveränderungen bei Experimentellem Rift-Valley-Fieber und ihre Beziehungen zur gelbfieberpathologie," Arquiv Insti Biol (Sao Paulo), 11:215-220, 1940. (In German).
60. Schulz, K.C.A.: "The Pathology of Rift Valley Fever or Enzootic Hepatitis in South Africa," J S African Vet Med Assoc, 22:3:113-120, 1951.
61. Mims, C.A.C.: "Rift Valley Fever Virus in Mice. VI. Histological Changes in the Liver in Relation to Virus Multiplication," Australian J Exptl Biol Med Sci, 35:6:595-604, 1957.
62. Mims, C.A.C.: "Viruses as Pathogenic Agents," Bull Epiz Dis Africa, 4:316-317, 1956.
63. Mims, C.A.C.: "The Coagulation Defect in Rift Valley Fever and Yellow Fever Virus Infections," Ann Trop Med Parasitol, 50:2:147-149, 1956.
64. Stefanopoulo, G.J., and Nagano, Y.: "Essais de Serothérapie Contre la Fievre de la Vallée du Rift ou Hepatite Enzootique," Rev Pathol Comparee et Hyg Gen, 38:1169-1176, 1938. (In French).

1. The first part of the report deals with the general situation of the country and the progress of the work during the year.

2. The second part contains a detailed account of the work done in each of the various departments.

3. The third part gives a summary of the results of the work and a statement of the financial position.

4. The fourth part contains a list of the names of the persons who have been employed during the year.

5. The fifth part contains a list of the names of the persons who have been elected to the various offices.

6. The sixth part contains a list of the names of the persons who have been elected to the various offices.


7. The seventh part contains a list of the names of the persons who have been elected to the various offices.

8. The eighth part contains a list of the names of the persons who have been elected to the various offices.


9. The ninth part contains a list of the names of the persons who have been elected to the various offices.

10. The tenth part contains a list of the names of the persons who have been elected to the various offices.

65. MacKenzie, R.D.: "Immunization of Mice Against Rift Valley Fever," J Pathol Bacteriol, 40:1:65-73, 1935.
66. Daubney, R., and Hudson, J.R.: "Rift Valley Fever," Lancet, 1:611-612, 1932.
67. Findlay, G.M., and Howard, E.M.: "The Effects of Cortisone and Adrenocorticotrophic Hormone on Poliomyelitis and on Other Virus Infections," J Pharm Pharmacol, 4:1:37-42, 1952.
68. Findlay, G.M., and MacCallum, F.C.: "Chemotherapy of Virus Diseases (correspondence)," Brit Med J, 1:875, 1938.
69. Phillips, A.P.: "Potential Antivirals. I. Simple Analog of Chloramphenicol (Chloromycetin)," J Amer Chem Soc, 74:6125-6127, 1952.
70. Greenhalgh, N.; Hull, R.; and Hurst, E.W.: "The Antiviral Activity of Acridines in Eastern Equine Encephalomyelitis, Rift Valley Fever, and Psittacosis in Mice and Lymphogranuloma Venereum in Chick Embryos," Brit J Pharmacol, 11:2:220-224, 1956.
71. MacKenzie, R.D., and Findlay, G.M.: "The Production of a Neurotropic Strain of Rift Valley Fever Virus," Lancet, 230:5864:140-141, 1936.
72. Findlay, G.M., and MacKenzie, R.D.: "Studies on Neurotropic Virus: Spontaneous Encephalomyelitis in Mice," Brit J Exptl Pathol, 17:441-447, 1936.
73. Naude, W. du T.; Madsen, T.; and Polson, A.: "Different Sized Infective Particles of Rift Valley Fever Virus," Nature, 173:1051-1052, 1954.
74. Polson, A.: "Weight Relationship Among Animal Viruses," Nature, 172:1154-1155, 1953.
75. Polson, A., and Linder, A.M.: "Determination of Sedimentation Constants of Protein and Viruses with the Help of the Spinco Preparative Ultracentrifuge," Biochem et Biophys Acta, 11:199-208, 1953.
76. Polson, A., and Madsen, T.I.: "A Brain Factor Influencing the Viability of Neurotropic Rift Valley Fever," Nature, 176:645-646, 1 October 1955.
77. Craig, J.F.: Fleming's Veterinary Obstetrics, 4th Edition, Bailliere, Tindall and Cox, London, England, 1930.
78. Saddington, R.S.: "In vitro and in vivo Cultivation of the Virus of Rift Valley Fever," Proc Soc Exptl Biol Med, 31:6:693-694, 1934.



79. MacKenzie, R.D.: "The Cultivation of the Virus of Rift Valley Fever," J Pathol Bacteriol, 37:1:75-79, 1933.
80. Endo, M.: "Sur le Variant Neurotrope de Virus de la Fievre de la Vallee du Rift Obtenu par la Culture du Tissue," Viruses, 1:42-50, 1951. (In Japanese).
81. Takemori, N.; Nakano, M.; and Hemmi, M.: "Plaque Formation with Rift Valley Fever Virus," Virology, 1:2:250-251, 1955.
82. Takemori, N.; Nakano, M.; Hemmi, M.; and Kitaoka, M.: "Propagation of Rift Valley Fever Virus in Ascites Hepatoma Cells of the Rat: Production of a New Variant of the Virus," Virology, 1:1:58-82, 1955.
83. Takemori, N.; Nakano, M.; Hemmi, M.; Ikeda, H.; Yanagida, S.; and Kitaoka, M.: "Destruction of Tumor Cells by Rift Valley Fever Virus," Nature, 174:698-700, 1954.
84. Iwasa, S.: "Multiplication of Rift Valley Fever Virus in Human Liver Cell Culture with Special Reference to Production Complement-Fixing Antigen," Japan J Exptl Med, 29:4:323-334, 1959.
85. Plowright, W., and Ferris, R.D.: "Rift Valley Fever. Tissue Culture," E African Vet Res Organization, Ann Rpt, 28-29, 1957.
86. Mims, C.A.C.: "Rift Valley Fever Virus in Mice. II. Adsorption and Multiplication of Virus," Brit J Exptl Pathol, 37:2:110-119, 1956.
87. Matumoto, M.; Ogiwara, H.; and Skinkawa, E.: "Multiplication of Neurotropic Rift Valley Fever Virus in Ehrlich Ascites Tumor Cells," Japan J Exptl Med, 25:6:255-265, 1955.
88. Naude, W.D., and Polson, A.: "Interference Between Active and Ultra-violet-Radiated Rift Valley Fever Virus," J Gen Microbiol, 16:2:491-497, 1957.
89. Joya, K.: "Antigenic Power of the Virus of Rift Valley Fever Inactivated by Ultraviolet Irradiation," Japan J Bacteriol, 7:6:613-616, 1952. (In Japanese).
90. Scott, G.R., and Whitcomb, M.A.: "Rinderpest Virus in Laboratory Animals: Interference of Rift Valley Fever Virus by Rinderpest Virus," E African Vet Res Organization, Ann Rpt, 16, 1956-1957.
91. Findlay, G.M., and MacCallum, F.O.: "An Interference Phenomenon in Relation to Yellow Fever and other Viruses," J Pathol Bacteriol, 44:2:405-424, 1937.



92. Isaacs, A.: "Measuring Concentration of Animal Viruses," Advances in Virus Research, IV. Academic Press, Inc., N.Y., N.Y., 1957.
93. Mims, C.A.C.: "Rift Valley Fever Virus in Mice, IV. Incomplete Virus; its Production and Properties," Brit J Exptl Pathol, 37:2:129-143, 1956.
94. Mims, C.A.C., and Mason, P.J.: "Rift Valley Fever Virus in Mice. V. The Properties of a Hemagglutinin Present in Infective Serum," Brit J Exptl Pathol, 37:5:423-433, 1956.
95. Andrewes, C.H., and Horstmann, D.M.: "The Susceptibility of Viruses to Ethyl Ether," J Gen Microbiol, 3:290-297, 1949.
96. Endo, M.: "A propos de la Conservation des Virus dans le Liquide d'oeuf," Japan J Exptl Med, 20:817-822, 1950. (In French).
97. Polson, A., and Madsen, T.I.: "A Brain Factor Influencing Viability of Neurotropic Rift Valley Fever," Nature, 178:4483:645-646, 1955.
98. Mims, C.A.C.: "Rift Valley Fever Virus in Mice. III. Further Quantitative Features of the Infective Process," Brit J Exptl Pathol, 37:2:120-128, 1956.
99. Findlay, G.M.: "The Transmission of Louping Ill to Monkeys," Brit J Exptl Pathol, 13:3:230-236, 1932.
100. Findlay, G.M.: "The Relation Between Dengue and Rift Valley Fever," Trans Roy Soc Trop Med Hyg, 26:2:157-160, 1932.
101. Broom, J.C., and Findlay, G.M.: "Complement-Fixation in Rift Valley Fever," Lancet, 1:222:609-611, 19 March 1932.
102. Casals, J.: "Viruses: Versatile Parasites. I. Arthropod-Borne Animal Viruses," Trans N Y Acad Sci, 19:219-235, 1957.
103. Bauer, D.J., and Bradley, P.L.: "The Xanthine Oxidase Groups. A Phenomenon Associated with the Multiplication of Neurotropic Viruses," Brit J Exptl Pathol, 37:5:447-460, 1956.
104. Gear, J.H.S.: "Rift Valley Fever in South Africa," Letter to the Editor, S African Med J, 25:35:620, 1951.

REFERENCES

- Adamson, J.S.: "Report of Director of Veterinary Services for Year Ending 30 September 1952, Southern Rhodesia," pp 18 and 29 (RVF), 1952.
- Anonymous: "Viruses and Infection Through the Skin," Brit Med J, 4216: 587, 1941.
- Araujo, E. de.: "Em Torno da Febre do Vale do Rift (Rift Valley Fever)," Cult Med, 2:113-119, 1932.
- Assumpcao, L.: "Relacoes Imunitarias Entre Febre Amarela e Outras Infeccoes Clinicamente Parecidas Que se Observam Nos Climats Tropicais e Sub-tropicais," Arquiv Hig Saude Publ, 9:19-32, 1944.
- Bauer, D.J.: "The Anti-Viral and Synergic Actions of Isatin Thiosemicarbazone and Certain Phenoxypyridimides in Vaccinia Infection in Mice," Brit J Exptl Pathol, 36:1:105-114, 1955.
- Camara, N.J.G. da: "O Aparecimento da 'Rift Valley Fever' na Uniao de Africa do Sol," Gas Do Agri, 3:115-116, 1951. (In Italian).
- Camara, N.J.G. da: "O Aparecimento da 'Rift Valley Fever' na Uniao da African do Sol," Ren Med Vet (Lisbon), 46:338:227-229, 1951.
- Cambessedes, H.: "Fievre de la Vallee du Rift," Rev Med Hyg Trop, 28:55-56, 1936.
- Culver, E.H.: "Rift Valley Fever," S African Inst Med Res, Ann Rpt, 38-39, 1951.
- Curasson, G.: "La 'Fievre de la Vallee du Rift' existe-t-elle au Soudan Francais?," Bull Soc Pathol Exotique, 27:6:599-602, 1934. (In French).
- Davis, D.H.S.: "Studies in Arthropod-Borne Viruses of Tongaland. III. The Small Wild Mammals in Relation to the Virus Studies," S African J Med Sci, 22:55-61, 1957.
- Editorial Board: "Status of Names of Bacterial Genera that are Later Homonyms of Names of Protozoan Genera," Intern Bull Bact Nomen and Tax, 3:2/3:109-110, 1953.
- Gledhill, A.W.: "Some Veterinary Diseases of Medical Interest," Brit Med Bull, 9:237-241, 1953.
- Henning, M.W.: Textbook. Animal Diseases in South Africa, 3rd Edition, Central News Agency, Johannesburg, South Africa, 1956.



- Horning, E.S., and Findlay, G.M.: "II. Microincineration Studies of the Liver in Rift Valley Fever," J Roy Microscop Soc, 54:1-9, 1934.
- Hurst, E.W.; Peters, J.M.; and Melvin, P.: "The Action of Mepacrine and Tryptan Red in a Number of Virus Diseases," Brit J Pharmacol, 7:473-481, 1952.
- Kaschula, V.R.: "Rift Valley Fever," Proc Reg Mtgs on Foreign Animal Dis, Rutgers University, New Brunswick, N.J., December 1956. pp. 113-118.
- Kaschula, V.R.: "Rift Valley Fever as a Veterinary and Medical Problem," J Amer Vet Med Assoc, 131:5:219-221, 1957.
- Koseki, Y.: "Bio- and Histochemical Studies on Mouse Liver Infected with Virus of Rift Valley Fever. I. Histochemical and Manometrical Studies," Sapporo Med J, 8:5/6:261-269, 1955. (In Japanese, English Summary).
- Koseki, Y.: "Bio- and Histochemical Studies on Mouse Liver Infected with Virus of Rift Valley Fever. II. Biochemical Studies," Sapporo Med J, 9:1; 1956. (In Japanese, English Summary).
- Lacorte, J.G.: "Virus Pantropics. IV. O Virus da Febre de Vale Rift," Rev Brasil Med, 7:7:461-464, 1950.
- Matumoto, M.; Iwasa, S.; and Endo, M.: "Complement-Fixation Reaction of Rift Valley Fever Virus," Japan J Exptl Med, 20:4:501-508, 1950.
- Matumoto, M.; Saburi, Y.; and Nishi, I.: "Rift Valley Fever Virus in the One-Day-Old Chick Embryo," J Immunol, 82:3:219-225, 1959.
- Matumoto, M.; Nishi, Y.; and Saburi, Y.: "Conditions Necessary for the Interference of the Neurotropic Virus with the Pantropic Virus of Rift Valley Fever," Compt Rend Soc Biol, 153:10:1645-1648, Feb 1960. (In French).
- Matumoto, M.; Nishi, I.; and Saburi, Y.: "Interference in Rift Valley Fever. Influence of the Neurotropic Strain on Multiplication of the Pantropic Strain," Compt Rend Soc Biol, 153:10:1649-1692, February 1960. (In French).
- McKercher, D.G.; Biberstein, E.L.; and Wada, E.M.: "A Review of Recent Findings in Infectious Disease of Sheep. I. Virus Diseases," J Amer Vet Med Assoc, 549-555, 1959.
- Nagano, Y.: "Investigations Immunologiques du Virus de la Fievre de Rift Valley," Japan Med J, 1:14-17, 1948. (In French).

- Nagano, Y.: "Virus-Serum Vaccination Contre la Fievre de la Vallee du Rift," Acta Convent Tert de Trop, 1:688-690, 1938. (In French).
- Nagano, Y.; Sawa, I.; Furano, S.; and Funabashi, T.: "Influence du Virus Inactive sur l'Infection par le Meme Virus," Japan J Exptl Med, 20:3:401-407, 1949. (In French).
- Nagano, Y., and Sonoda, T.: "Vaccination Contre la Fievre de la Vallee du Rift Avec le Virus Inactive par la Chaleur," Ren Pathol Comparee et Hyg Gen, 40:16-21, 1940.
- Najera, L.E.: "Active Immunization and Viruses; Modern Aspects of Vaccination Against Rift Valley Fever, American Macular Fever, and Q Fever," Ann Med Pub, 4:581-596, 1952.
- Nosengo, C.: "Della probabile Esistenza Della Febre Della Valle Del Rift," Prog Vet, 3:60-63, 1948. (In Italian).
- Notani, N.: "Studies on Effects of Activities of Mouse Liver Fractionatants by Rift Valley Fever Virus Infection. I. Study on Fractionating Procedures. II. Distribution of Virus Particles in Fractionated Cell Components," Sapporo Med J, 11:4:215-227, 1957.
- Pellissier, A.: "Enquete Serologique sur l'incidence des Virus Neurotropes on Afrique Equatoriale Francaise," Bull Soc Pathol Exotique, 47:2:223-227, 1954.
- Pellissier, A., and Rousselot, R.: "Enquete serologique sur l'incidence des Virus Neurotropes ches Quelques Signas de l'Afrique Equatoriale Francaise," Bull Soc Pathol Exotique, 47:2:228-231, 1954.
- Sardou, M.: "La Fievre de la Vallee du Rift," Gas med de France, 674-681, 1935.
- Sawa, I.: "Inhibition de la Multiplication du Virus de la Fievre de la Vallee du Rift par le Virus Homologue Irradie par des Rayons Ultra-violet," Compt Rend Soc Biol, 149:21:2050-2052, 1955. (In French).
- Sawa, I.: "Inhibition de Multiplication du Virus de la Fievre de la Vallee du Rift par le Virus Homologue Irradie par des Rayons Ultra-violet, II.," Compt Rend Soc Biol, 150:835-837, 1955. (In French).
- Sawa, I.; Shibuki, M.; and Takeuti, S.: "A propos de la Duree Minima Necessaire d/incubation de Melange de Virus et d'immunserum dans la Reaction de Neutralization," Japan J Exptl Med, 20:689-697, 1950. (In French).
- Scott, G.R., and Hirsch, R.B.: "Rift Valley Fever and Rift Valley Rodents," E African Med J, 36:665-667, December 1959.

- Scott, G.R.; Weddell, W.; and Reid, D.: "Preliminary Findings on the Prevalence of Rift Valley Fever in Kenya Cattle," Bull Epiz dis African, 4:17-25, 1956.
- Shahan, M.S., and Traum, J.: "The Exotic Zoonoses," Ann N Y Acad Sci, 70:3:614-623, 1958.
- Sosov, R.F.: "Rift Valley Fever (Enzootic Hepatitis)," Veterinariia, 32:12:69-71, 1955.
- Stefanopoulo, G.J.: "Sur le 'diounde' a propos d'une Enquete Epidemiologique sur le Fievre Jaune dans les Pays de Segou et de Macina (Soudan, Francais)," Bull Soc Path Exot, 26:4:560-562, 1933.
- Stefanopoulo, G.J., and Nagano, Y.: "Sur la Vaccination contre La Fievre de la Vallee du Rift," Rev Pathol Comparee et Hyg Gen, 38:1297-1301, 1938.
- Theiler, Max: "Rift Valley Fever," in Viral and Rickettsial Infections of Man, 3rd Edition, T. Rivers, Editor, J.B. Lippincott, Co., pp. 391-394, 1959.
- Van Rooyen, C.E., and Rhodes, A.J.: Virus Diseases of Man, 2nd Edition, Thomas Nelson and Sons, New York, 1948.
- Verge, J.: "Le Maladies Communes a l'Homme and Aux Animaux. IV. Une Entite Exotique Nounelle; Le Fievre de la Valle du Rift," Ren Gen Med Vet, 41:485-272-277, 1932.
- Weinbren, M.P., and Mason, P.J.: "Rift Valley Fever in Arnicanthis abyssinicus," E African Virus Inst Rpt, 7:11, 1956.



JUN 18 2008

OCT 16 2008



<http://nihlibrary.nih.gov>

10 Center Drive
Bethesda, MD 20892-1150
301-496-1080

NIH LIBRARY



3 1496 00995 3129